Summary of the Data Elements for

Each Landscape Position

The following tables provide the range of values for selected attributes within each landscape position (NSLP). Two analyses are provided. The first column shows the field attribute (column or row heading) as found in the NSLP table. The second column gives the range of values found in the database for that attribute within that landscape position. The datum followed by the percent of the total area it represents within that category is provided. The third column gives the range of values found in the database for that attribute within that landscape position. The datum followed by the percent of the map units it represents within that category is provided. The following are descriptions of the attribute fields provided in the analysis:

1STSURTEX

This field contains the surface soil texture code for the Sequence Number 1 soil series found within the map unit. Elements include:

C = clay

FS= fine sand

FSL= fine sandy loam

GR-MUCK= gravelly muck

GRV-L= very gravelly loam

LFS= loamy fine sand

LS= loamy sand;

MARL= marl

MUCK= muck

MK-FS= mucky fine sand

MK-LFS= mucky loamy fine sand

MK-S= mucky sand

ROCK= rock

S = sand

SCL= sandy clay loam

SICL= silty clay loam

SIL= silt loam.

ECOCOM

This field contains the number of the ecological community as defined by the Natural Resource Conservation Service's (formerly the Soil Conservation Service) publication "The 26 Ecological Communities of Florida. Elements are:

NA= no data

#1= North Florida Coastal Strand

#2= South Florida Coastal Strand

#3= Sand Scrub

#4= Longleaf Pine-Turkey Oak Hills

#6= South Florida Flatwoods

#8= Cabbage Palm Flatwoods

#9= Everglades Flatwoods

#10= Cutthroat Seeps

#11= Upland Hardwood Hammocks

#12= Wetland Hardwood Hammocks

#13= Cabbage Palm Hammocks

#14= Tropical Hammocks

#15= Oak Hammocks

#16= Scrub Cypress

#17= Cypress Swamp

#18= Salt Marsh

#19= Mangrove Swamp

#21= Swamp Hardwoods

#22= Shrub Bogs- Bay Swamps

#24= Sawgrass Marsh

#25= Freshwater Marsh

#26= Slough.

HYDRIC

This indicates if a COMPNAME is defined as a Hydric soil. If the COMPNAME is hydric the entry is "Y". If the COMPNAME is not hydric the entry is "N".

SHWT RANGE

This is the high and low value for the average range of the seasonal high water table. Units are in feet.

SHWT DURATION

This is the average month in which the seasonal high water table begins and ends.

HYDROGRP

This is the Hydrologic Group for a particular COMPNAME. The codes used with a general description of the soil conditions associated with them are as follows:

A= High infiltration rates. Soils are deep, well drained to excessively drained sands and gravels.

A/D= Drained/undrained hydrology class of soils that can be drained and are classified.

B= Moderate infiltration rates. Deep and moderately deep, moderately well and well drained soils that have moderately coarse textures.

B/D= Drained/undrained hydrology class of soils that can be drained and are classified..

C= Slow infiltration rates. Soils with layers impeding downward movement of water, or soils that have moderately fine or fine textures.

C/D= Drained/undrained hydrology class of soils that can be drained and classified

D= Very slow infiltration rates. Soils are clayey, have a high water table, or are shallow to an impervious layer.

DRAINCLASS

This is the soil drainage class code, which identifies natural drainage condition of the soil. The code refers to the frequency and duration of periods when the soil is free of saturation. These codes are defined as follows:

E= excessively

MW= moderately well

P= poorly

SE= somewhat excessively

SP= somewhat poorly

VP= very poorly

W= well.

SOILRUN

This indicates the potential of a COMPNAME to allow chemicals to leave the application site with runoff water and/or detached soil particles. Possible entries are LOW, MEDIUM, or HIGH. Additional information concerning soilrun and other types of Water Quality Interpretations is in Section II-iii-L of the Field Office Technical Guide (FOTG) published by the Florida NRCS.

Distribution of Data Within NSLP Categories

NSLP Category 2: Tidal Soils

Summary By Area	Summary By Map Units
NIΛ	MUCK (400/) FC (460/)

1STSURTEX NA MUCK (40%), FS (16%), S (16%), MARL

(8%), SIL (6%)

ECOCOM NA (2%), #19 (70%), #18 (28%) NA (16%), #19 (53%), #18 (31%)

HYDRIC YES (100%) YES (100%)

SHWT Range (ft) 0.0 to 0.5 (94%), 1.0 (4%), 0.0 to 6.0 0.0 to 0.5 (84%), 0.0 to 6.0 (16%)

(2%), Other (less than 1%)

SHWT Duration JAN to DEC (100%), Other (less than JAN to DEC (96%), JUN to OCT (2%), JUN to

SEP (2%)

HYDROGRP D (100%) D (100%)

DRAINCLASS VP: (94%), P: (6%) P: (32%), VP: (68%) **RUNOFF** HIGH (100%) HIGH (100%)

Major Soil Series Durbin & Wulfert (31%), Wulfert Beaches (16%), Kesson (16%), Perrine (6%),

(11%), Peckish (9%), Kesson (7%), Durbin AND Wulfert (4%)

Key Largo (5%), Estero (4%)

Area (non-water) 96591 HA 96591 HA

NSLP Category 3: Marl and Rocky Soils

	Summary By Area	Summary By Map Units
1STSURTEX	NA	SIL (31%), MARL (25%), FSL (13%), ROCK
		(13%), SICL (13%)
ECOCOM	#25 (59%), #24 (36%), #26 (4%),	#24 (44%), #25 (32%), #14 (6%), #18 (6%),
	#14 (1%), #18 (less than 1%)	#26 (6%), ND (6%)
HYDRIC	YES: (100%, excluding rock areas)	YES: (100%, excluding rock areas)
SHWT Range (ft)	0.0 to 1.0 (64%), -1.0 to 1.0 (35%),	0.0 to 1.0 (62%), -1.0 to 1.0 (19%), 0.5 to 1.0
	0.5 to 1.0 (less than 1%), 0.0 to 0.5	(6%), 0.0 to 0.5 (13)
	(less than 1%)	, ,
SHWT Duration	JUN to SEP (38%), JUN to OCT	JUN to SEP (31%), JUN to OCT (13%), JUN
	(17%), JUN to NOV (31%), JUN to	to NOV (50%), JUN to DEC (6%)
	DEC (14%)	
HYDROGRP	D (100%)	D (100%)
DRAINCLASS	P (78%), VP (21%), SP (1%)	P (81%), VP (13%), SP (6%)
S-RUNOFF	H (74%), M (26%)	HIGH (75%), MED (25%)
Major Soil Series	Biscayne (38%), Perrine (21%),	Pennsuco (25%), Perrine (19%), Biscayne
-	Ochopee (17%), Rock Outcrop -	(16%), Ochopee (13%)
	Vizcaya - Biscayne Complex (14%)	. , . , ,
Area (non-water)		86574 HA

NSLP Category 4: Everglades Peat

	Summary By Area	Summary By Map Units
RTEX	NA	MUCK (100%)

1STSURTEX NA MUCK (100%) **ECOCOM** #25 (40%), #24 (60%) #24 (56%), #25 (44%)

HYDRIC YES (100%) YES (100%)

SHWT Range (ft) -2.0 to 0.0 (68%), -1 to 0.0 (32%) -2.0 to 0.0 (78%), -1 to 0.0 (22%) **SHWT Duration** JUN to APR (68%), JUN to FEB (22%)

(32%)

 HYDROGRP
 D (100%)
 D (100%)

 DRAINCLASS
 VP (100%)
 VP (100%)

 S-RUNOFF
 HIGH (100%)
 HIGH (100%)

Major Soil Series Pahokee (32%), Terra Ceia (31%), Dania (28%), Lauderhill (28%), Pahokee

Lauderhill (22%), Dania (9%) Muck (22%), Terra Ceia Muck (17%), Torry

Muck (6%)

Area (non-water) 297470 HA 297470 HA

NSLP Category 5: Muck Depressions

ary By Map Units
ì

1STSURTEX NA FS (1%), LFS (1%), MUCK (98%)

ECOCOM #25 (78%), #24 (4%), #21 (5%), #19 ND (1%), #25 (66%), #24 (13%), #22 (13%),

(3%), #17 (3%), Others (less than #21 (3%), #19 (1%), #17 (3%), #12 (1%)

1%)

HYDRIC Y (100%) Y (100%)

SHWT Range (ft) -2.0 to 0.0 (70%), -1.0 to 0.0 (14%), - -2.0 to 0.0 (65%), -1.0 to 0.0 (16%), -2.0 to

2.0 to 1.0 (5%), 0.0 to 0.5 (9%), -1.0 1.0 (4%), 0.0 to 0.5 (9%), -1.0 to 1.0 (5%), 0.0

to 1.0 (2%), 0.0 to 1.0 (1%) to 1.0 (1%)

SHWT Duration JUN to OCT (11%), JUN to DEC JUN to OCT (16%), JUN to DEC (6%), JUN to

(3%), JUN to JAN (7%), JUN to FEB JAN (8%), JUN to FEB (1%), JUN to MAR (1%), JUN to MAR (1%), JUN to APR (5%), JUN to APR (57%), JAN to DEC (6%)

(72%), JAN to DEC (6%)

 HYDROGRP
 D (100%)
 D (100%)

 DRAINCLASS
 VP (100%)
 VP (100%)

S-RUNOFF HIGH (100%) HIGH (99%), MED (1%)

Major Soil Series Samsula (17%), Hontoon (13%), Gator (10%), Samsula (10%), Terra Ceia

Kaliga (11%), Gator (8%), Okeelanta (10%), Hontoon (9%), Okeelanta (9%),

(7%), Sanibel (7%), Terra Ceia (7%) Sanibel (9%)

Area (non-water) 266548 HA 266548 HA

NSLP Category 6: Sand Depressions

Summary By Area Summary By Map Units

1STSURTEX C (1%), FS (59%), FSL (6%), LFS (4), LS (2), NA

MK-FS (9%), MK-LFS (2), MK-S (1%), S

23(16%), SCL (1%)

ECOCOM NA (2%), #26 (6%), #25 (81%), #21 NA (1%), #12 (4%), #16 (1%), #17 (6%), #21 (9%), #16 (2%)

(14%), #22 (2%), #25 (69%), #26 (4%)

HYDRIC YES (100%) YES (100%)

SHWT Range (ft) -2.0 to 0.0 (67%), -2.0 to 1.0 (19%), -2.0 to 0.0 (70%), -2.0 to 1.0 (9%), 0.0 to 0.5

> 0.0 to 0.5 (7%), 0.0 to 1.0 (7%), (9%), 0.0 to 1.0 (11%), Others (1%)

Others (less than 1%)

SHWT Duration JUN to SEP (17%), JUN to OCT JUN to SEP (11%), JUN to OCT (9%), JUN to

> (7%), JUN to NOV (1%), JUN to DEC NOV (3%), JUN to DEC (4%), JUN to JAN (3%), JUN to JAN (2%), JUN to FEB (2%), JUN to FEB (4%), JUN to MAR (63%),

(11%), JUN to MAR (58%), JUN to JUN to APR (1%), JUL to MAR (3%)

APR (1%).

HYDROGRP D (100%) D (100%)

DRAINCLASS VP (92%), P (8%) VP (89%), P (11%) S-RUNOFF HIGH (100%) HIGH (99%), MED (1%)

Major Soil Series Riviera (13%), Basinger (10%), Floridana (9%), Chobee (9%), Felda (7%),

Boca, Riviera, Limestone Substratum Basinger (6%)

and Copeland (9%), Floridana (7%),

Chobee (5%), Felda (5%)

Area (non-water) 534906 HA 534906 HA

NSLP Category 7: Flats Soils

Summary By Area **Summary By Map Units**

1STSURTEX NA FS (81%), LFS (1%), LS (1%), MK-FS (1%),

S (15%), SCL (1%)

ECOCOM NA (2%), #9 (4%), #8 (2%), #6 (2%), NA (1%), #6 (1%), #8 (%), #9 (1%), #10 (2%),

#26 (61%), #25 (8%), #22 (2%), #16 #11 (1%), #12 (3%), #13 (4%), #14 (1%), #16

(2%), #14 (2%), #13 (4%), #12 (2%), (1%), #22 (1%), #25 (3%), #26 (78%)

#11 (2%), #10 (4%), #26 (4%)

HYDRIC NO (3%), YES (97%)

SHWT Range (ft) 0.0 to 1.0 (94%), 0.0 to 0.5 (2%), -1.0 0.0 to 1.0 (84%), 0.0 to 0.5 (14%), -1.0 to 1.0

to 1.0 (4%)

SHWT Duration JUN to SEP (8%), JUN to OCT (8%), JUN to SEP (9%), JUN to OCT (22%), JUN to

> JUN to NOV (41%), JUN to DEC NOV (37%), JUN to DEC (9%), JUN to FEB

(13%), JUN to FEB (25%), JUN to (15%), JUN to MAR (7%)

MAR (5%)

HYDROGRP D (100%) D (100%)

DRAINCLASS P (100%), VP (Less than 1%) P (99%), VP (1%) S-RUNOFF HIGH (100%), MED (Less than 1%) HIGH (99%), MED (1%)

Major Soil Series Basinger (20%), Pineda (16%), Basinger (12%), Pineda (13%), Pompano

> Riviera (12%), Malabar (10%), (9%), Malabar (8%)

Holopaw (7%), Valkaria (7%)

713013 HA Area (non-water) 713013 HA

NSLP Category 8: Flatwood Soils

Summary By Area Summary By Map Units

1STSURTEX NA FS (72%), LFS (1%), LS (1%), S (26%) **ECOCOM** #9 (1%), #8 (3%), #6 (74%), #15 #6 (84%), #8 (5%), #9 (1%), #12 (5%), #13

(4%), # 13 (9%), #12 (4%), #11 (5%)

(1%), #6 (3%)

HYDRIC NA NO (98%), YES (2%)

SHWT Range (ft) 0.5 to 1.5 (99%), Others (less than 0.5 to 1.5 (99%), Others (1%)

1%)

SHWT Duration JUN to SEP (93%), JUN to OCT JUN to SEP (90%), JUN to OCT (2%), JUN to

(2%), JUN to NOV (1%), JUN to FEB NOV (3%), JUN to FEB (5%)

(4%)

 HYDROGRP
 D (99%), C (1%)
 C (1%), D (99%)

 DRAINCLASS
 P (95%), SP (5%)
 P (96%), SP (4%)

 S-RUNOFF
 HIGH (100 %)
 HIGH (100%)

Major Soil Series Immokalee (20%), Myakka (16%), Oldsmar (10%), Wabasso (10%), Immokalee

Smyrna (13%), Oldsmar (8%), (9%), Myakka (8%), Hallandale (6%)

Wabasso (7%)

Area (non-water) 1473609 HA 1473609 HA

NSLP Category 9: Knolls

Summary By Area	Summary By Map Units
-----------------	----------------------

1STSURTEX NA FS (59%), GR-MUCK (1%), GRV-L (3%),

MUCK (2%), S (34%), SICL (1%)

ECOCOM NA (9%), #9 (9%), #6 (10%), #4 NA (9%), #1 (1%), #11 (8%), #14 (3%), #15

(10%), #3 (34%), #24 (2%), #2 (9%), (4%), #2 (6%), #24 (1%), #3 (38%), #4 (12%),

#15 (2%), #14 (5%), #11 (8%), #1 #6 (14%), #9 (5%)

(2%)

HYDRIC N (100%) N (100%)

SHWT Range (ft) 1.5 to 3.5 (16%), 2.0 to 3.5 (30%), 2.0 to 3.5 (30%), 1.5 to 3.5 (16%), 3.5 to 6.0

3.5 to 6.0 (26%), 1.5 to 3.0 (2%), 3.5 (12%), 1.5 to 3.0 (11%), 3.0 to 5.0 (8%), 5.0 to 5.0 (3%), 4.0 to 5.0 (11%), 3.0 to 5.0 (7%), 3.5 to 5.0 (4%), 4.0 to 5.0 (4%), 5.0 (2%), 4.0 to 6.0 (2%), 5.0 to 6.0 (4%), 2.0 to 3.0 (3%), 2.0 to 8.0 (2%), 1.0 to 3.0 (5%), Others (less (2%), 2.5 to 3.5 (2%), 2.5 to 5.0 (2%), 0.0 to

than 1%) 6.0 (1%), 1.0 to 3.0 (1%)

SHWT Duration JUN to OCT (4%), JUN to NOV JUN to OCT (10%), JUN to NOV (39%), JUN

(46%), JUN to DÉC (21%), JUN to to DEC (12%), JUN to FEB (1%), JUL to OCT FEB (3%), JUL to OCT (6%), JUL to (13%), JUL to NOV (15%), JUL to JAN (3%), NOV (15%), JUL to JAN (2%), JAN JAN to DEC (2%), AUG to FEB (1%), JUN to

to DEC (1%), AUG to FEB (1%), SEP (4%), JUL to DEC (2%), Others (less than 1%)

HYDROGRP C (52%), B (14%), A (34%) A (28%), B (11%), C (61%)

DRAINCLASS MW (47%), MW-P (2%), SE (3%), MW (46%), MW-P (1%), SE (2%), SP (4%),

SP (36%), SP-MW (2%), W (10%) SP-MW (1%), W (7%)

S-RUNOFF HIGH (14%), LOW (50%), MED HIGH (8%), LOW (40%), MED (52%)

(36%)

Major Soil Series Pomello (16%), Tavares (16%), Pomello (13%), Tavares (5%), Satellite (8%),

Krome (10%), Zolfo (10%), Satellite Archbold (5%), Adamsville (6%), Canaveral (9%), Archbold (5%), Chekika (5%), (5%), Electra (5%), St. Augustine (5%),

Sparr (4%)

Area (non-water) 242235 HA 242235 HA

NSLP Category 10: Dune and Central Ridge

Summary By Area Summary By Map Units

1STSURTEX NA FS (46%), S (54%)

ECOCOM NA (3%), #4 (28%), #3 (45%), #2 NA (1%), #11 (5%), #2 (13%), #3 (51%), #4

(17%), #11 (7) (28%) N (100%) N (100

SHWT Range (ft) 6.0 to 6.0 (98%), 0.0 to 6.0 (2%)

SHWT Duration N/A (98%), JUN to OCT (2%)

HYDROGRP A (100%)

DRAINCLASS E (92%), W (6%), Others (2%)

S-RUNOFF LOW (99%), MED (1%)

Major Soil Series Candler (45%), Astatula (24%),

Neilhurst (7%), St. Lucie (5%)

Area (non-water) 135398 HA

HYDRIC

N (100%)

6.0 to 6.0 (99%), 0.0 to 6.0 (1%) N/A (99%), JUN to OCT (1%)

A (100%)

E (79%), E-P (1%), W (18%)

LOW (95%), MED (5%)

Astatula (10%), Candler (18%), Paola (15%),

St. Lucie (21%)

135398 HA